

**UPAVISHA KUCHALA (STRYCHNOUS NUXVOMICA LINN) – A REVIEW****Chauhan Mitesh<sup>1</sup>, Chavhan Kalpana<sup>2</sup>, Suryawanshi Shama<sup>3</sup>**<sup>1</sup>PG Scholar, <sup>2</sup>Assistant Professor, <sup>3</sup>Professor Department of Agadtantra, Government Ayurveda College, Nagpur, Maharashtra, India**ABSTRACT**

*Kuchala (strychnous nuxvomica Linn)* is a well known poisonous plant in Indian system of medicine. It is included in *Upavisha* by ayurveda texts. Strychnine is a popular folk medicine from ancient times. Even today also many people in ruler India use *Kuchala* in medicine form. According to *Charaka* even an acute poison can become an excellent drug if it is properly administered, and similarly even a drug, if not properly administered, becomes an acute poison. *Kuchala* is a known vegetable poison to Ayurveda as well as modern science but it is used in many medicinal preparations of Ayurveda and other allied medicinal pathies. Strychnine is a main contain of *kuchala* which was first used medically in 1540, and continued to be used in many stimulants, Tonics and cathartics. This review article includes the overall information about the poisonous plant *Kuchala*, it's, Toxicological aspect, Medico-legal aspect and therapeutic uses mentioned in Ayurveda and in other systems of medicine.

**Keywords:** *Kuchala, strychnous nuxvomica, Upavisha, Therapeutic uses, spinal poisons*

**INTRODUCTION**

*Kuchala (strychnous nuxvomica Linn)*, a well known plant in Indian system of medicine is being used extensively in different classical formulations with great therapeutic significance. It has been stated categorically that strong poisons could be the best medicine, if it is used after proper detoxification (*shodhana*), in proper therapeutic dose and formulation. On the contrary, a good medicine may affect adversely unless it is used for proper person in proper dose <sup>[1]</sup>. *Rasratnasamucchaya* described eleven number of *Upavisha* <sup>[2]</sup>. In Ayurvedic literature *Upavisha* are the group of drugs which were less toxic in nature and not so lethal but produce certain toxic symptoms on consumption or administration. They are having less toxic potency <sup>[3]</sup>. Though the plant *Kuchala* is described under the '*Upavisha varga*' (sub poisonous group) it's seeds have been used

successfully in different formulations to combat different diseases after proper *Shodhan sanskar* (processing of purification). Strychnine is a main content of *Kuchala* is popular in folk medicine from ancient period. *Nuxvomica* was introduced in Europe in the sixteenth century, but was not used in medicine. This alkaloid strychnine has been in use as a rodenticide at that time. It is sometime used for killing stray dogs hence the name dog buttons is used for it, being chiefly employed to poison dogs, cats, crows etc. Strychnine was first used medically in 1540 and continued to be used in many stimulants Tonics and cathartics until as recently as the 1960s <sup>[4]</sup>. The seeds are mainly used as Aphrodisiac, Appetizer, Anti-periodic, Digestive, Purgative, and Stimulant. It's also used in Anemia, Asthma, Bronchitis, and Intermittent and malarial fever. Physicians of

ayurveda successfully employed this drug & preparations containing it in a number of diseases after proper purification. Many Ayurveda medicinal formulations like Agnitundirasa, Laxmivilasarasa, Shulnir-mulanarasa, Suptivaatarirasa, Vishatinduka<sup>[5]</sup>. contains Kuchala beej as their basic ingredient. This review article is a sincere

attempt to summarize the information concerning about poisonous drug Kuchala (strychnous nuxvomica Linn) described in Indian system of medicine in respect to its literary, pharmacological activity, toxicological effects, and therapeutic uses in various systems of medicines including Ayurveda.



**Scientific Classification**<sup>[6]</sup>:

<b>Kingdom:</b>	<b>Plantae</b>
<b>(unranked):</b>	Angiosperms
<b>(unranked):</b>	Eudicots
<b>(unranked):</b>	Asterids
<b>Order:</b>	Gentianales
<b>Family:</b>	Loganiaceae
<b>Genus:</b>	<i>Strychnos</i>
<b>Species:</b>	<i>S. nuxvomica</i>

**Botanical Name**– *Strychnous nuxvomica*

**Family**–Loganiaceous, Karaskara Kula

**Vernacular Names**<sup>[7]</sup>

**Hindi Name**- Kuchala

**English Name**- Nuxvomica

**Telugu Name**- Mushini Ginjala, Mushti Vittulu

**Bengali Name**- Kunchila

**Marathi Name** -Kajara

**Gujarati Name**- Jherkuchala, Zerkochala

**Tamil Name**- Yettikottai

**Malayalam Name**- Kaajjeel

**Arabian Name** – Ajaraki, Habbul Gurav

**Parsi Name** – Kuchula, Phuloosemaahi

**Sanskrit Synonyms:**<sup>[8]</sup> *Kuchelaka, kuchel,*

*Kuchila, Kuchil, Vishatinduk, Tindu, Tinduk, Vishatinduk, Karaskara, Ramyafala, Kupaak, Vishamushtika, Vishamushti, Kaalkuta.*

**Classification:**

**Ayurveda:** *Sthavara Vanaspatik vish Upavisha*<sup>[9]</sup> *phala visha (beeja visha)*<sup>[10]</sup>

**Modern medicine:** Neurotoxin spinal excitant poison<sup>[11]</sup>

**Distribution:**<sup>[12]</sup>

It is found throughout tropical India up to an altitude of 360 m, in Uttar Pradesh, Bihar, Orissa, Coromandel Coast, Andhra Pradesh and Karnataka. It is most common in the forests along the western coasts.

**Plant Description:**<sup>[13]</sup>

Plant is dense, hard white and close-grained. The branches are irregular and are covered with a smooth ashen bark. The young shoots are deep green color with a shiny coat. The leaves have an opposite arrangement, short stalked, are oval shaped, also have shiny coat and are smooth on both sides. The leaves are about 4 inches (10cm) long and 3 inches (7.6cm) wide. The flowers are small with a pale green color with a funnel shape. They

bloom in the cold season with a smooth and have a foul smell. The fruit are about the size of a large apple with a smooth and hard shell which when ripened is a lovely orange color. The meat of the fruit is soft and white with a jelly like pulp containing five seeds covered with a soft woolly substance. The seeds are removed from the fruit when ripe. They are then cleaned, dried and sorted. The seeds have the shape of flattened disk completely covered with hairs radiating from the centre of the sides. This gives the seeds very characteristics sheen. The seeds are very hard, with a dark gray horny endosperm where the small embryo is housed that give off no odor but possess a very bitter taste. The plant is native to south East Asia and Australia normally in tropical and subtropical areas.

**Major chemical constituents:** <sup>[14]</sup>

- Brucine
- Strychnine
- Vomisine
- Kajine and Novocain(N-methyl pseudobrucine)
- Strychnine and Isostrychnine
- Cuchiloside
- Loganic acids

**Ayurveda view of Kuchala-** In Bruhat Trayi(3 basic granthas of Ayurveda i.e. *Charaka Samhita, Sushrut Samhita and Vagbhata Samhita*) and *Dhanvantari Nighantu* did not mention *Vishatinduka* or *kupillu*. *Shodhala* denoted it as *Visha Tinduk* and included it in *karveeradi varga* <sup>[15]</sup>, while *Bhavamishra* described it as *Kakattinduka* or *Kupilu* <sup>[16]</sup>. *Kaideva Nighantu* quated a drug *Vishamusti*, which may be *nuxvomica* <sup>[17]</sup>. *Rajanighantu* described *Kuchala* in *Prabhadradi varga* <sup>[18]</sup>. He also quated five types of *Vishamushti*. In ethnomedicine other species of the same genus are in vogue in Telangana (a state in India). These species are used in the name

of different kinds of '*Mushini*'. In modern era, due to its poisonous nature, *Nuxvomica* was very reluctantly introduced into the European pharmacopoeias.

**Ayurvedic properties** <sup>[19]</sup>

**Rasa:** -*Katu Tikta Guna*:-*Rruksha, Laghu, Teekshna, Veerya*:-*Ushna, Vipaka*:- *Katu*

**Doshaghanata:** *Kaphavatshamak* <sup>[20]</sup>  
*Kaphapittanashanam* <sup>[21]</sup>

**Rogghnata [22]:** *Sandhivata, Amavata, Vrana, Kushatha, Nadishoola, Ardhangha, Gatibhransha, Gyanabhransna, Peshiposha, Kampa, Badhirya, Ardita, Pakshaghata, Andria, Amadya, Amashyastha, Amadosha, Grahani, Udarshoola, Arsha, Krimi, Rakta-vikara, Vatarakta, Hridyashaithilaya, Hridayodara, Kasa, Phuphusshotha, Dhwa-jabhangha, Sheeghrapatana, Daurbalya, Kushtha, Kandu, Atisweda, Vishamajwara, Visuchika.*

**Karma:** *Shothahara, Puthihara, Vedanasthapana, Uttejaka, Nadibalya, Deepana, Pachana, Grahi, Shoolprashamana, Hridayottejaka, Kaphaghna, Kasahara, Vajikarna, Balya, Katupaushtika, Kushthaghna, Kandughana, Swedapnayana.*

**Kupilu – uses –** <sup>[23]</sup>

*Hanti Meda* – lowers cholesterol, useful in obesity

*Krumihara* – useful in intestinal worm infestation

*Shvasahara* – useful in asthma and wheezing

*Gulmahara* – useful in abdominal tumor, bloating

*Arshohara* – useful in hemorrhoids

*Mushikavishahara* – useful in rat bite

*Vishtambhi* – causes constipation

*Rochana* – improves taste, useful in anorexia

*Agnikrut* – improves digestion strength

*Grahi* – absorbent, useful in diarrhea

*Kushtahara* – useful in skin disorders

*Pramehajit* – useful in urinary disorders, diabetes

*Madakrut* – causes intoxication

*Kanthamayahara* – useful in diseases of throat

**Kupilu Shodhana (Detoxification / Purification method)** <sup>[24]</sup>

1. Fry *Kuchala* seeds with ghee in a pan on slow flame till it's outer covering become led-yellow coloured. Take these seeds and remove the outer skin of seeds and grind the hot pulp immediately. This *shodhana* process is useful in emergency use of *Kuchala*.
2. Wrap *Kuchala* seeds in a cloth, keep it in *Dolayantra* with cow's milk, and boil it for 3 hrs. After 3 hrs remove the seeds, grind it in iron *Kharal*, and use the *churna* (powder). Skin of seeds is removed. It is boiled with milk for 7 days, dried, then it is fried in ghee and powdered.

**3. Medicinal Dose:-** 1/2 to 1 *Gunja* <sup>[25]</sup>

**Ayurvedic preparations of Kupilu:** - <sup>[26]</sup> *Navjeevan Rasa, Agnitundi rasa, Laxmivilas Rasa, Shoolanirmulan Rasa, Suptivatri Rasa, Sarameha Vishapaha Yoga, Vishatinduk Taila* (External use).

**Modern view of kuchala:-** <sup>[27]</sup> Strychnine was first used medically in 1540 and continued to be used in many stimulants, tonics and cathartics until as recently as the 1960s <sup>[28]</sup>. *Kuchala* was recognized in America (1830) before it was official in England, although long before this date it was a dispensatory drug. In 1799, however, it was official in the pharmacopeia bo-russica, and in two other continental European pharmacopeias. The dominating constituents of nuxvomica is a complex compound which in natural form, is an invaluable remedy, by means of chemical reagents it can be split into parts, embracing two intensely poisonous alkaloid products, a glucosidal and acids. These alkaloids are Strychnine, Brucine, and perhaps Igasurine (yet in doubt). The main acid is Igasuric

acid, while the glycoside is named loganin. These are all colorless bodies, the alkaloids being very bitter, and energetically poisonous, Brucine being a poison similar to strychnine, acting with less violence and more slowly, but not less surely, than strychnine.

**Strychnine:**  $C_{21}H_{22}N_2O_2$  this violently poisonous alkaloid is crystalline, slightly soluble in cold water the solution being alkaline and bitter. Strychnine is said to be the bitterest substance in the world. The taste is detectable even in a dilution of 1/100,000 or more. It dissolves in 7 with parts of chloroform and 150 parts of 90% alcohol. Strychnine is a terrible titanic poison, affecting the cerebro-spinal system, but it kills without producing marked anatomical changes, the muscles and nerves being scarcely altered, although brain and spinal cord may be congested, stomach and limbs intensely congested, right side of heart gorged (sometimes empty) and the lungs congested, the fatal dosed of strychnine is as low as 1/2 grain. Indeed, it is recorded that 1/16 grain killed a two year old child in four hours, while 1/2 grain killed a man in twenty minutes.

**Brucine:** this related alkaloid is also a product of chemical action on nuxvomica. It is known to chemists as dimethoxyl-strychnine

**Brucine-**  $C_{22}H_{26}N_2O_4$ . is very bitter feebly soluble in cold alcohol. It differs in reaction from strychnine in that strong Sulphuric or Nitric acid strikes with it blood – red color, whereas with strychnine no coloration appears. Brucine is a poison which has the physiological, but in markedly less degree. Authorities differ, some considering it one sixteenth; others from one – fortieth to one fiftieth less energetic than strychnine as convulstant. The antidotes and treatment for poisoning by Brucine are the same as for strychnine.

### Pharmaceutical preparations:

The alkaloids are sometimes prescribed in solution and have long, made into pill, tablet, or pellet form of the solutions, half solution of strychnine is an old preparation occasionally used at present.

**Tincture of nuxvomica:** This preparation was among the first official preparation of *Kuchala*. It is made today by dissolving 20 grams of dried extract of nuxvomica in enough mixture of alcohol 3 volumes, volume to make 1000cubic centimeters. This is perhaps the best known drug store preparation of nuxvomica and needs no special comment.

**Mode of Action** <sup>[29]</sup>:- **Strychnine** stimulates all parts of the CNS and particularly the anterior horn cells of spinal cord causing greatly increased reflex excitability. Normal inhibition of motor cell stimulation is lost so that any slight stimulus such noise, light, or air breeze causes violent generalized muscle spasms.

### Pharmacological Activities <sup>[30]</sup>

Anti HIV, Hepatoprotective, Anticholestatic, Ant lipid Peroxidative property, Antiulcer, Insecticidal, CNS stimulant and Strychnine showed remarkable negative chronoscopic activity on frog isolated heart and guinea pig atria and activity retained in vivo also (open chest dog). Strychnine (50mg/kg) when injected subcutaneously increased levels of acetylcholine in spinal cord and sustained convulsions in frog for 4 hr. Isostrychnine N-oxide and isobrucinne N- oxide showed the most potent cytotoxicity to tumour cell lines of K562, HELA & HEP-2.

**Part used** <sup>[31]</sup>- Seed is the most used part of this herb. Rarely, root bark is also used. It should be purified before using for medicinal purposes.

**Dosage** <sup>[32]</sup>- Seed powder –A possible fatal dose is ½ gm [ below 33 mg].

**Toxic symptoms** <sup>[33]</sup> –

- Bitter taste
- Twitching and stiffness of muscles of face and neck
- Convulsions- initially clonic i.e. intermittent and then tonic i.e. sustained.
- Any stimulus like movements of patient, noise, touch, light or water immediately produces convulsions.
- Muscles became rigid and stiff, so that body is thrown in to the form of arch
- Synosis
- Blood stained froth at nose and mouth
- Eyes :-prominent and staring, with dilated pupils
- Mind remains clear till end
- Death is painful

### Diagnosis of Poisoning <sup>[34]</sup>

- TLC gives reliable qualitative results on gastric aspirate, urine, blood or tissues.
- Best specimens are urine and gastric aspirates
- HPTLC provides accurate quantitative data.
- Blood levels in the range of 0.1 to 0.3 mg/100ml are generally lethal.

### Post Mortem Appearance:-

- Rigid attitudes characteristic of the clinical state may persist for a long time after death. <sup>[35]</sup>
- There may be oozing and hemorrhages are usually present in muscles.
- As in death following any violent muscular activity, the lymph in thoracic duct is bloody.
- The spasm of the muscles interferes with respiration and causes death from asphyxia.
- Early onset and disappearance of rigor mortis.
- Postmortem calorcity
- Dilated pupils.

### Forensic Significance of Plant <sup>[36]</sup>:-

- Strychnine has been uncommonly employed in murder owing to various obvious reasons like bitter taste, dramatic nature of symptoms-that will always arouse suspicion of foul play, and easy delectability in body fluids and tissues.
- Accidental poisoning can result in children who chew on the seeds out of curiosity while playing or foraging in the countryside.
- Previously, therapeutic misadventures used to be fairly common when strychnine was an approved constituent of various over-the-counter tonics and cathartics.
- Accidental poisoning can also result from inadvertent consumption of strychnine-containing rodenticide.
- Owing to the agonizing nature of death, strychnine is rarely employed in suicide.

#### **Action and Uses** <sup>[37]</sup>

The root is bitter, tonic, febrifuge and useful in cholera, intermittent fever and bites of venomous reptiles. The leaves are applied as poultice in the treatment of chronic wounds and ulcers and leaf decoction is useful in paralytic complaints. The pulp of the ripe fruit is used in treating paralytic affections of palms and foot. The seeds are bitter, nerving, tonic, Alexiteric, Aphrodisiac, Appetizer, Ant periodic, Antihelminthic, Emetic, Digestive, Purgative, Diabetes, Colic, Intermittent And Malarial Fever, Insomnia, Cardio spasms, Skin Diseases, Nerve Debility, Dyspepsia, Diarrhea, Dysentery, Hysteria, Mental Emotions, Epilepsy, Chronic Constipations, Gout, Chronic Rheumatism, Hydrophobia, Spermatorrhoea, Opium or Lead poisoning, Paralysis and weakness of limbs. The wood is used in Dysentery, Dyspepsia and Fevers.

*Strychnous nuxvomica* is also used in homeopathy.<sup>[38]</sup> It is said in Homeopathy "if you do not know what should be prescribed, then give Nux Vomica." It is often used as an antidote for over drugging. Nux Vomica is generally prescribed for males who are thin, irritable and lose temper by slight provocation. Also, for those who do a good deal of mental work, study a lot or handle business affairs and lead indoor life. Because of mental strain, such people often seek the help of stimulants, such as coffee, liquor, or use sedatives like opium or any other cannabis preparation. People, who take rich food, attend parties and generally overindulge themselves until late at night, often have irregular bowel movements (or have constipation). They often take laxatives like Hajmola, liver tonics, etc. Nuxvomica soothes and calms overexcited nervous system and improves digestion and bowel movement. It increases Appetite, vigor and gives potency to males who have ruined themselves by excessive use of stimulants. That is why it is called the medicine of "bigrey Nawab (spoilt men). It may be taken in low potency of 6 or 30 (in case of irritable, overexcited persons), and above 200 or more potency in case of habitually constipated and hard drinkers. It is one of the best remedy for mania-a-potu (acute alcoholism). Nux Vomica of 6 and 30 potency should be taken once a day before going to sleep at night. If it is 200 potency then it should be taken once a week. If it is still higher, then once a fortnight <sup>[39]</sup>

#### **DISCUSSION**

*Kuchala* is a well known spinal poison to modern science. It is used in Ayurvedic pharmacopeia from ancient period. Ayurveda texts like *Rasatarangini*, *Rasratnasamucchaya*, *Raj-Nighantu*, and *Bhavprakash* mentioned detail description of the plant, basic properties, therapeutic uses,

medicinal preparations. Some Ayurveda texts like *Bruhat- Trayi* (3 basic granthas of Ayurveda i.e. *Charaka Samhita*, *Sushruta Samhita* and *Vagbhata Samhita*) and *Dhanvantari Nighantu* did not mention *Kuchala*. Even in *Kalpasthanu Sushruta* described types of *visha* according to *adhithana* (a part of plant were poison resides), among it he includes *fala visha* (poisonous fruits), but he didn't mention in it. Due to some properties like *Ashukaritwa*, *Ushna*, *Teekshna vish dravya* get spread rapidly in the body. So for the quick action of medicines many *Ayurvedic* formulations contain these *vishadravyas* like *Kuchala* as their ingredient. By utilizing these properties of *vishadravyas* medicines can be made more effective. So we found that many *Rasashastra* based texts are having description of poisonous drugs like *Kuchala* in detail. Rastarangini stated the detoxification process of *Kuchala*, so that purified *Kuchala* can get used in medicinal formulations. Modern toxicology includes it in a deadly poison. It is categorized as Neurotoxin spinal excitant poison. Medico legally this plant is important too. Homicidal Death due to *Kuchala* is uncommon because of bitter taste, dramatic symptoms and easy detectability in body fluids and tissues. Accidental poisoning is common among children. Homeopathy also mentions many therapeutic uses of *Kuchala*. In homeopathic material medica *Nuxvomica* is mentioned as laxative, Digestive, Increasing vigor and vitality in male and also useful in alcoholism.

## CONCLUSION

*Kuchala (strychnous nuxvomica Linn)* is one of the deadly poisons known to mankind. Though it is poison, it is important part of Ayurvedic and Homeopathy pharmacopeia. It is a basic ingredient of many ayurveda formulations. Due to properties like *Ashukaritwa*, *Ushna*, *Teekshna vish*

*dravya* like *Kuchala* get spread rapidly in the body. So for the quick action they are used in medicinal formulations of Indian system of medicine and other systems.

## REFERENCES

1. Agnivesh, Charak Samhita, edited by Vaidya Yadavji Trikamji Acharya, Chaukhambha surabharti Prakashan, Varanasi 2008.
2. Shastri Ambikadatta (Suratnojiyala Hindi Commentry), Rasaratnasammuchaya, Chukhambha Amarbharti Prakashan, Varanasi, Ed. 8<sup>th</sup>, 1988, Peg no.170.
3. Dr.Namburi Shekhar U.R, A Textbook Of Agadtantra, Chukhambha Sanskrit Sansthan, Varanasi, Reprint 2013, peg no. 16.
4. Pillay V.V., Comprehensive Medical Toxicology, 2<sup>nd</sup> Ed, Paras Medical Publisher, Hyderabad, India, peg no.878
5. Rasatarangini edited by Pandit Kashinath Shasti, Motilal Banarasidas, 41 U.A. Banglo Road, Jawahar Nagar, Delhi 110007, 11th edition
6. [https://en.wikipedia.org/wiki/Strychnous\\_nuxvomica](https://en.wikipedia.org/wiki/Strychnous_nuxvomica).
7. Dr.Shastry J.L.N., Dravyaguna vijjana, Vol-2, Chukhambha Orientaliya, Varanasi, Ed. 3<sup>rd</sup>, 2008, peg no.353.
8. Pandit Kashinath Shastry, Rasatarangini, Motilala Banarasidas, Delhi, Ed. 11<sup>th</sup>, 1979, peg no 676.
9. Pandit Kashinath Shastry, Rasatarangini, Motilala Banarasidas, Delhi, Ed. 11<sup>th</sup>, 1979, peg no 675.
10. Shastri Ambikadatta, editors, Sushrut-samhita-klp.3/3, Chukhambha Sanskrit Sansthan, Varanasi, Reprint 2000, Peg No .16.
11. Singhal S.K., Toxicology At A Glance, The National Book Depot, Mumbai, Ed, 7th, reprint 2009, Peg. No.114.

12. Prof.Lavekar G.S., Database on Medicinal Plants Used in Ayurveda and Siddha, Vol-5, CCRAS,New Delhi, Reprint 2008,peg no.139.
13. Trivedi K.P., Dhanvantari Vanaushadhi Visheshanka (Hindi), Dhanvantari karyalalya,Vijayghar,Vol 2,peg no 248-259.
14. Dr.Shastry J.L.N.,Dravyaguna vijna- na,Vol 2,Chukhambha Orientaliya, va- ranasi, Ed.3<sup>rd</sup>,2008,peg no.353.
15. .Sharma P.V., Shodhal Nighantu, Oriental Institute, Baroda, 1st Ed. 1978, peg no.125.
16. Mishara B. and Vaishya R., Bhavapra- kasha,Purwardhwam,Chukhambha Sanskrit sansthana, Varanasi, Ed.8, 1993, peg no.568.
17. Dr.Shastry J.L.N.,Dravyaguna vijna- na,Vol 2,Chukhambha Orientaliya, va- ranasi, Ed.3<sup>rd</sup>, 2008, peg no.353.
18. Tripathi indradev, Raj-Nighantu, Krushnadas academy, Varanasi, 1st Ed., 1982, peg no 293.
19. Prof.Lavekar G.S., Database on Me- dicinal Plants Used in Ayurveda and Siddha, Vol-5, CCRAS, New Delhi, Reprint 2008, peg no.140.
20. Sharma P.V.,Shodhal Nighantu, Orient- al ,Institute, Baroda,1<sup>st</sup> Ed. 1978, Peg no.125.
21. Mishara B.and Vaishya R., Bhava- prakasha, Purwardhwam, Chukhambha Sanskrit sansthana, Varanasi, Ed.8,1993, peg no.568.
22. Sharma P.,Dravyaguna-Vigyana,@nd part,Chukhambha Bharti Academy, Varanasi,Reprint 2005,peg.No.85
23. <http://easyayurveda.com/2014/01/08/kupilu-nux-vomica-uses-dose-purification-side-effects/>
24. Pandit Kashinath Shastry, Rasataran- gini, Motilala Banarasidas, Delhi, Ed, 11,1979, peg no 679.
25. Yadav AcharyaTrikamji, Dravyaguna Vigyanam, 2<sup>nd</sup>Part, Sharma ayurved Mandir, Datiya,Ed. 5<sup>th</sup>2001,peg no. 270.
26. Pandit Kashinath Shastry, Rasataran- gini, Motilala Banarasidas, Delhi ,Ed, 11,1979,peg no 684-688.
27. John Uri Lloyd,Drug Treatise Number 8,Issued by Lloyd Brothers, Cincinati, Ohio.,www.swsbm.com
28. Pillay V.V., Comprehensive Medical Toxicology, 2<sup>nd</sup> Ed,Paras Medical Pub- lisher,Hydrabad,India,pegno.876.
29. Dr.Parikh C.K., Textbook of Medical Jurisprudence Forensic Medicine and Toxicology,CBS Publishers and Dis- tributors, NewDelhi, Ed.6<sup>th</sup>, reprint 2007, peg no.10.57.
30. Prof.Lavekar G.S., Database on Medi- cinal Plants Used in Ayurveda and Siddha,Vol-5,CCRAS,NewDelhi ,Reprint 2008, peg no.140.
31. Dr.Shastry J.L.N.,Dravyaguna vijna- na,Vol 2,Chukhambha Orientaliya, va- ranasi, Ed.3<sup>rd</sup>,2008,peg no.354.
32. Pryce D.M. And Ross C.F.,Ross's Post Mortem Apperances,Oxford University Press, New York, Ed.6<sup>th</sup> 1963, peg .no.38.
33. Singhal S.K.,Toxicology At A Glance,The National Book Depot, Mumbai, Ed, 7<sup>th</sup>, Reprint 2009, Peg. no.114
34. Pillay V.V., Comprehensive Medical Toxicology,2<sup>nd</sup> Ed,Paras Medical Pub- lisher,Hydrabad,India,pegno.877.
35. Pryce D.M. And Ross C.F.,Ross's Post Mortem Apperances,Oxford University Press,New York,Ed.6<sup>th</sup> 1963, peg.no. 38.
36. Pillay V.V., Comprehensive Medical Toxicology, 2<sup>nd</sup> Ed, Paras Medical Publisher, Hydrabad, India, peg no.878.



37. Prof.Lavekar G.S., Database on Medicinal Plants Used in Ayurveda and Siddha, Vol-5, CCRAS, NewDelhi, Reprint 2008, peg no.140.
38. Roland Hofbauer, Eva Pasching, Doris Moser & Michael Frass (2010). "Heparin-binding epidermal growth factor expression in KATO-III cells after *Helicobacter pylori* stimulation under the influence of *Strychnous nuxvomica* and *Calendulaofficinalis*". *Homeopathy* **99** (3): 177–182.
39. Homeopathic Journal :: Volume: 5, Issue: 2, Dec 2011 (new Papers) from Homeorizon.com

---

### **CORRESPONDING AUTHOR**

**Dr. Chauhan Mitesh**

PG Scholar, Department of Agadtantra, Government Ayurveda College, Nagpur, Maharashtra, India

**Email:** cmitesh306@gmail.com

---

*Source of support: Nil*  
*Conflict of interest: None Declared*