# INTERNATIONAL AYURVEDIC MEDICAL JOURNAL



International Ayurvedic Medical Journal, (ISSN: 2320 5091) (November, 2017) 5(11)

# CRITICAL STUDY OF STHAULYA (OBESITY) AND ITS PRINCIPLES OF MANAGEMENT

#### Pranit Hanumantrao Patil

Assistant Professor, R.J.V.S., Bhaisaheb Sawant, Ayurvedic Medical College, Sawantwadi, Dist. Sindhudurg, Maharashtra, India

Email: pranit.patil2950@gmail.com

#### **ABSTRACT**

Obesity is a medical condition in which excess body fat has accumulated to the extent that it may have a negative effect on health. Obesity increases the likelihood of various diseases and conditions, particularly Cardiovascular diseases, Respiratory Diseases, Type 2 diabetes, obstructive sleep apnea, certain types of cancer, osteoarthritis and depression. Obesity is more common in women than men. Authorities view it as one of the most serious public health problems of the 21st century. Obesity is stigmatized in much of the modern world (particularly in the Western world), though it was seen as a symbol of wealth and fertility at other times in history and still is in some parts of the world. In 2013, the American Medical Association classified obesity as a disease. In India, the prevalence of overweight increased from 9.7% near the turn of the century to nearly 20% in studies reported after 2010. For children and adolescents, these studies show that obesity and overweight are rapidly increasing, not just in the higher income groups but also in the rural poor, where under nutrition and underweight remain major health concerns. Here the word 'Critical' means clear and impartial. In present study an effort made to find etiopathogenesis, impact of *Sthaulya* and management of Obesity critically which will provide baseline data to future researchers regarding management of obesity?

Keywords: Sthaulya, Obesity Treatment, Ayurvedic, Shodhana, Santarpana Vyayama

#### INTRODUCTION

Obesity is the complex, multifactorial type of nutritional metabolic disease. It occurs due to imbalance between intake of energy and its expenditure resulting in positive energy balance, which is characterized by the abnormal

growth of the adipose tissue, resulting in an increase in the body weight to the extent of 20% or more of the standard weight for the person's age, sex and height. Obesity is a medical condition in which excess body fat

has accumulated to the extent that it may have a negative effect on health. Obesity increases the likelihood of various diseases and conditions, particularly Cardiovascular diseases, Respiratory Diseases, Type 2 diabetes, obstructive sleep apnea, certain types of cancer, osteoarthritis and depression. Obesity is a leading preventable cause of death worldwide, with increasing rates in adults and children. In 2015, 600 million adults (12%) and 100 million children were obese. Obesity is more common in women than men. Authorities view it as one of the most serious public health problems of the 21st century. Obesity is stigmatized in much of the modern world (particularly in the Western world), though it was seen as a symbol of wealth and fertility at other times in history and still is in some parts of the world. In 2013, the American Medical Association classified obesity as a disease [1]. In India, the prevalence of overweight increased from 9.7% near the turn of the century to nearly 20% in studies reported after 2010. For children and adolescents, these studies show that obesity and overweight are rapidly increasing, not just in the higher income groups but also in the rural poor, where under nutrition and underweight remain major health concerns [2]. However, India has also seen a surge in obesity. It had 0.4 million obese men, or 1.3% of the global obese population in 1975, but in 2014, it zoomed into the fifth position with 9.8 million obese men, or 3.7% of the global population. Among women, India has jumped to the third rank with 20 million obese women (5.3% of global population) [3]. Here the word 'Critical' means clear and impartial. In present study an effort made to find etiopathogenesis, impact of Sthaulya and management of Obesity critically. Obesity is a

state of excess adipose tissue mass. The most widely used method to gauge obesity is the body mass index (B.M.I.), which is equal to weight/height2 (in kg/m2). Other approaches to quantifying obesity include anthropometry (skin -fold thickness), densitometry (underwater weighing), computed tomography (CT) or magnetic resonance imaging (MRI), and electrical impedance. Based on data of substantial morbidity, a BMI of 30 is most commonly used as a threshold for obesity in both men and women [4].

#### MATERIALS AND METHODS

Critical study of Sthaulya is categorized under

- **A)** Etymology
- **B)** Definition of *Sthaulya*
- C) Nidana
- **D)** Rupa
- E) Impact of Sthaulya
- F) Samprapti
- **G)** Management of *Sthaulya*

#### A) ETYMOLOGY

- The word *Sthaulya* is delivered from root "*Sthu*" with suffix "*Ach*", which stands probably for thick or solid or strong or big or bulky.
- The term *Sthaulya* is derived from the *AtmanepadiSthoolayate Dhatu* having meaning to grow, increase <sup>[5]</sup>
- The word Obesity is derived from the Latin 'obesitas', which means "stout, fat, or plump" [6]

## B) DEFINITION OF STHAULYA

MedomamsaativruddhatvatChalaSphig Udara Stanaha |

 $Ay a thop a chay ot saho Naro a tis thool Uchya te \\ ||^{[7]}$ 

The person is called over obese who, due to excessive increase of fat and muscles, has pendulous buttocks, abdomen and breasts, and suffers from deficient metabolism and energy.

## C) *NIDANA* [8]

All the causative factors described in *Ayurvedic* classics can be classified under 5 groups

- 1) Aharatmaka Nidana
- 2) Viharatmaka Nidana
- 3) Manas Nidana
- 4) Beeja Dosha
- 5) Anya Nidana

# 1. AHARATMAKA HETU (DIETERY FACTORS)

Factors related to improper consumption of diet

Atisampurana(Over eating), Santarpana, Adhyasana

- Factor related to quality of food material Guru Aharasevana, MadhuraAharasevana, SheetaAharasevana, SnigdhaAharasevana, Sleshmala Aharasevana,
- Specific food substances causing obesity
  Navannasevana, Nava Madhya Sevana,
  GramyaRasasevana, Audak Rasasevana,
  Mamsa Sevana, PayaVikar Sevana, Dadhi
  Sevana, Sarpi Sevana, IkshuVikara Sevana,
  Guda Vikara Sevana, Shali Sevana, Godhum
  Sevana, Masha Sevana, Rasayan Sevana,
  VrushyaAaharaSevana

#### 2. VIHARATMAKA HETU

Avyavaya, Divaswapna, Asana Sukha, Swapnaprasangat, GandhamalaSevana, Bhojanotar Snanaare ViharakmakaHetu

# 3. MANSIKA HETU (PSYCHOLOGICAL FACTORS)

Harashnityatvat, Achintana, Priyadarshana, Manasonivritti are ManasjanyaHetu

#### 4. BEEJA DOSHA

Acharya Charaka has described a specific cause Beeja Dosha Swabhavafor Sthaulya, Acharya Vagbhata has described the Sahaja Sthaulya due to dietary fault of pregnant lady. Therefore, two miscellaneous causative factors of Sthaulya have been described in Ayurvedic text. i.e. Beejadoshajanya and Garbhajvyadhi.

#### 5. ANYA NIDANA

It includes *SnigdhaMadhuraBasti Sevana* (Administration of unctuous and sweet enema), *Tailabhyanga* (Massaging of oil), *SnigdhaUdvartana* (Unctuous action) etc.

# CRITICAL ANALYSIS OF *NIDANA* IN RELATION TO *STHAULYA*

## 1) Role of Aharatmaka Nidana in Sthulya

Ahararasa plays as major role for increasing Meda Dhatu in Sthaulya<sup>[9]</sup>. Sthaulya and Karshya depend upon the quality and quantity of Ahararasa. On the basis of Samanya VisheshSiddhanta, the excess food consumption of similar substance of similar quality, similar in action helps in the over production of Dhatu. In the same manner increased intake of Aharatmaka Nidana causes over production of Medodhatu.

# 2) Role of Viharatmaka Nidana in Sthaulya All the Aharatmaka Nidana ultimately decreases physical activity, which aggravates Kapha and leads of Meda deposition. Viharatmaka Nidana like Divaswapna having Abhishyandi property leads to blockage of the micro channels of the body, specifically in MedovahaStrotas.

#### 3) Role of Manasjanya Nidana in Sthaulya

According to *CharakacharyaHarshanity*a and *Achintana* are two psychological factors which are responsible for *MedaVriddhi*. These factors are *Kapha* aggravating factors lead of *Meda* deposition. With this type of psychological wellbeing and jolliness that person indulges more in worldly pleasure and excess energy stored in the form of *Meda*.

## 4) Role of Beejadosha in Sthaulya

Acharva Charaka has mentioned that Beejadosha major plays a MedoVruddhi. Defect in Beejabhagavayava, which resembles with Genes, may lead to defective development of that organ. Also, Bhavamishra has mentioned that increased proportionate of Meda and decreased proportion of Shukra in Beeja at the time of conception predisposes towards development of stout but weak body<sup>[10]</sup>. Moreover, over nutrition particularly with Madhura Rasa during pregnancy is mentioned as a causative factor for birth of obese child, which indicate role of hereditary factor in genesis of Sthaulya.

# CAUSES ACOORDING TO MODERN SCIENCE [11]

At an individual level, a combination of excessive food energy intake and a lack of physical activities are thought to explain most cases of obesity. A 2006 review identified ten other possible contributors to the recent increase of obesity: insufficient sleep, endocrine disruptors, decreased variability in ambient temperature, decreased rates of smoking, increased use of medications that can cause weight gain, proportional increases in ethnic and age groups that tend to be heavier, pregnancy at a later age, epigenetic risk factors passed on generationally, natural selection for

higher BMI, and assortative mating leading to increased concentration of obesity risk factors.

#### 1) DIET

A 2016 review supported excess food as the primary factor. Dietary energy supply per capita varies markedly between different regions and countries. The primary sources of these extra carbohydrates are sweetened beverages, which now account for almost 25 percent of daily food energy in young adults in America, and potato chips. Consumption of sweetened drinks such as soft drinks, fruit drinks, iced tea, and energy and vitamin water drinks is believed to be contributing to the rising rates of obesity and to an increased risk of metabolic syndrome and Type 2 diabetes. Vitamin D deficiency is related to diseases associated with obesity. As societies become increasingly reliant on energy-dense, big-portions, and fastfood meals, the association between fast-food consumption and obesity becomes more concerning.

#### 2) SEDENTARY LIFESTYLE

A sedentary lifestyle plays a significant role in obesity. Worldwide there has been a large shift towards less physically demanding work, and currently at least 30% of the world's population gets insufficient exercise. This is primarily due to increasing use of mechanized transportation and a greater prevalence of labor-saving technology in the home. In children, there appear to be declines in levels of physical activity due to less walking and physical education. In both children and adults, there is an association between television viewing time and the risk of obesity.

#### 3) GENETICS

Like many other medical conditions, obesity is the result of interplay between genetic and environmental factors. Polymorphisms in various genes controlling appetite and metabolism predispose to obesity when sufficient food energy is present. Studies that have focused on inheritance patterns rather than on specific genes have found that 80% of the offspring of two obese parents were also obese, in contrast to less than 10% of the offspring of two parents who were of normal weight. The thrifty gene hypothesis postulates that, due to dietary scarcity during human evolution, people are prone to obesity. Their ability to take advantage of rare periods of abundance by storing energy as fat would be advantageous during times of varying food availability, and individuals with greater adipose reserves would be more likely to survive famine. This tendency to store fat, however, would be maladaptive in societies with stable food supplies. This theory has received various criticisms, and other evolutionarily-based theories such as the drifty gene hypothesis and the thrifty phenotype hypothesis have also been proposed.

## 4) OTHER ILLNESSES AND MEDICA-TIONS

Certain physical and mental illnesses and the pharmaceutical substances used to treat them can increase risk of obesity. Medical illnesses that increase obesity risk include several rare genetic syndromes as well as some congenital or acquired conditions: hypothyroidism, Cushing's syndrome, growth hormone deficiency, and the eating disorders: binge eating disorder and night eating syndrome. Certain medications may cause weight gain or changes in body composition; these include insulin, sul-

fonylureas, thiazolidinediones, atypical antipsychotics, antidepressants, steroids, certain anticonvulsants (phenytoin and valproate), pizotifen, and some forms of hormonal contraception.

#### 5) SOCIAL DETERMINANTS

Many explanations have been put forth for associations between BMI and social class. It is thought that in developed countries, the wealthy are able to afford more nutritious food, they are under greater social pressure to remain slim, and have more opportunities along with greater expectations for physical fitness. In undeveloped countries the ability to afford food, high energy expenditure with physical labor, and cultural values favoring a larger body size are believed to contribute to the observed patterns. Malnutrition in early life is believed to play a role in the rising rates of obesity in the developing world. Endocrine changes that occur during periods of malnutrition may promote the storage of fat once more food energy becomes available.

#### 6) GUT BACTERIA

The study of the effect of infectious agents on metabolism is still in its early stages. Gut flora has been shown to differ between lean and obese humans. There is an indication that gut flora in obese and lean individuals can affect the metabolic potential. This apparent alteration of the metabolic potential is believed to confer a greater capacity to harvest energy contributing to obesity. Whether these differences are the direct cause or the result of obesity has yet to be determined unequivocally.

## D) *RUPA* SYMPTOMS OF *STHAULYA* [12]

Sweda (Excessive Ati Sweating), SharamjanyaShwasa (Breathlessness on mild exertion), AtiNindra (Excessive KaryaDorblyta (Difficulty to perform heavy work), Jadyatha (Sluggishness), Alpaayu (Short life span), Alpabala (Decreased bony strength), Uathashahani (Inertness), SharirDurgandhta (Foul odour of thebody), Gadgadtava (Unclear voice), Kshudhavridhi (Excessive hunger), AtiTrishna (Excessive thirst) are symptoms of Sthaulya.

# E) IMPACT OF *STHAULYA* COMPLICATIONS IN STHAULYA [13]

Obese individual may suffer from Jwara, Udara, Bhagandara, Prameha, Urustambha, Prameha Pitika, Urustambha etc disorders

# OBESITY RELATED ORGAN SYSTEMS REVIEW $^{[14]}$

Obesity can cause various disorders, possible system-wise enumeration enlisted below,

#### • Cardiovascular System

Hypertension, Congestive heart failure, Corpulmonale, varicose veins, pulmonaryembolism Coronary artery disease

#### Respiratory System

Dyspnea, Obstructive sleep apnea, Hypoventilation syndrome, Pickwickian syndrome, Asthma

## • Gastrointestinal System

Gastroesophageal reflux disease, Nonalcoholic fatty-liver disease, Cholelithiasis, Hernias, Colon cancer

#### • Endocrine System

Metabolic syndrome Type 2 diabetes Dyslipidemia Polycystic ovarian syndrome

#### Musculoskeletal System

Hyperuricemia and gout, Immobility, Osteoarthritis (knees and hips), Low back pain, Carpal tunnel syndrome

#### • Genitourinary System

Urinary stress incontinence, Obesity-related glomerulopathy, Hypogonadism (male)Breast and uterine cancer, Pregnancy complications

## • Psychological

Depression/low self-esteem, Body image disturbance, Social stigmatization

#### • Neurologic

Stroke, Idiopathic intracranial hypertension, Meralgiaparesthetica, Dementia

#### Integumentery system

Striaedistensae, Stasis pigmentation of legs, Lymphedema, Cellulitis, Intertrigo, carbuncles, Acanthosisnigricans, Acrochor dons (skin tags), Hidradenitissuppurativa

## F) SAMPRAPTI OF STHAULYA

In obesity, *Medo Dhatu* obstructs the channels related to digestive system (Koshta). This results in obstruction of Vata Dosha in the digestive system. It increases Agni (digestive fire) and dries up food in the stomach and intestines. Hence, food digestion speeds up increasing hunger. So the patient digests food quickly and becomes a voracious eater. If he does not get food on time, he becomes prone to serious disorders. The Agni (digestive fire), influenced by Pitta and Vata Dosha get vitiated. They burn the food as the forest fire burns the forest. Hence the body weight increases. Due to disproportionate increase of fat, diseases of very serious types are caused, all of a sudden, by Vata, etc. This may lead to instantaneous death. Owing to an excessive increase

of fat and muscle tissue, the buttock, abdomen and breast become pendulous and his strength is rendered disproportionate with his physical growth <sup>[15]</sup>.

In Samprapti of Sthaulya, all the three Doshas are vitiated especially Kledaka Kapha, Pachaka Pitta, Samana and Vyana Vayu are the responsible factors for proper digestion and metabolism of food at the level of alimentary tract and body tissue. Dushti of these Tri-Dosha components results in indigestion metabolic deformity and form action of Ama at tissue level as well as alimentary tract. Sthaulya is a Dushya dominant disorder. Due to excess Agni more Annarasais produced and

the quality of diet i.e. Guru, Madhur, Snigdha, SheetaGuna dominance Dhatu Poshakansha is formed in more quantity. At start the Rasagata, Raktagata and MamsagataSneha also increase production of Medodhatu. But due to Medodhatwagnimandyata, this condition worsens and nourishment of further Dhatus doesn't happen properly. Due to binge eating, Rasa, Rakta, MamsagataSneha tends to increase. Patient shows the symptoms of Rasavriddhi and Kaphavriddhi e.g. Angagaurav, Alasya, Nidradhikya. Medadhatu get increased with physical signs like ChalaSphik Udara Stana, KshudraShwas, Swedadhikva etc and finally lands into Sthaulya.

# Factors involved in Samprapti of Sthaulya [16]

• Doshas : Vata, Pitta, Kapha

• Dushyas : Rasa, Rakta, Mamsa, Meda

• Strotas : Rasavaha, Mamsavaha, Medavaha

• Strotodushti : Sanga

• Agni : TeekshnaJatharaagni

• Aam : Dhatwaagni-Maandya-Janya

UdbhavaSthana : Aamashaya VyaktaSthana : SarvaShareera

• Adhishthana : Mamsa Dhatu, Meda Dhatu

Rogamarga : BahyaRogamargaSadhyasadhyatva : KrucchraSadhya

# PATHOPHYSIOLOGY OF OBESITY [17]

There are many possible pathophysiological mechanisms involved in the development and maintenance of obesity. Investigators postulated that leptin was a satiety factor. In the ob/ob mouse, mutations in the leptin gene resulted in the obese phenotype opening the possibility of leptin therapy for human obesity. However, soon thereafter J. F. Caro's laboratory could

not detect any mutations in the leptin gene in humans with obesity. On the contrary Leptin expression was increased proposing the possibility of Leptin-resistance in human obesity. Since this discovery, many other hormonal mechanisms have been elucidated that participate in the regulation of appetite and food intake, storage patterns of adipose tissue, and development of insulin resistance. Since leptin's discovery, ghrelin, insulin, orexin, PYY 3-36, cholecystokinin, adiponectin, as well as many other mediators have been studied. The adipokines are mediators produced by adipose tissue; their action is thought to modify many obesity-related diseases. Leptin and ghrelin are considered to be complementary in their influence on appetite, with ghrelin produced by the stomach modulating short-term appetitive control. Leptin is produced by adipose tissue to signal fat storage reserves in the body, and mediates long-term appetitive controls.

# 1) GRAPHIC DEPICTION OF A LEPTIN MOLECULE

Leptin and Ghrelin are produced peripherally; they control appetite through their actions on the central nervous system. In particular, they and other appetite-related hormones act on the hypothalamus, a region of the brain central to the regulation of food intake and energy expenditure. The circuit begins with an area of the hypothalamus, the arcuate nucleus, that has outputs to the lateral hypothalamus (LH) and ventromedial hypothalamus (VMH), the brain's feeding and satiety centers, respectively. The arcuate nucleus contains two distinct groups of neurons. The first group co expresses neuropeptide Y (NPY) and agouti-related peptide (AgRP) and has stimulatory inputs to the LH and inhibitory inputs to the VMH. The second coexpressesprogroup opiomelanocortin (POMC) and cocaine- and amphetamine-regulated transcript (CART) and has stimulatory inputs to the VMH and inhibitory inputs to the LH. Consequently, NPY/AgRP neurons stimulate feeding and inhibit satiety, while POMC/CART neurons stimulate satiety and inhibit feeding. Both

groups of arcuate nucleus neurons are regulated in part by leptin. Leptin inhibits the NPY/AgRP group while stimulating the POMC/CART group. Thus a deficiency in leptin signaling, either via leptin deficiency or leptin resistance, leads to overfeeding or may account for some genetic and acquired forms of obesity.

# 2) PHYSIOLOGIC REGULATION OF ENERGY BALANCE<sup>[18]</sup>

Substantial evidence suggests that body weight is regulated by both endocrine and neural components that ultimately influence the effect or arms of energy intake and expenditure. This complex regulatory system is necessary because even small imbalances between energy intake and expenditure will ultimately have large effects on body weight. A major regulator of these adaptive responses is the adipocyte-derived hormone leptin, which acts through brain circuits to influence appetite, energy expenditure, and neuroendocrine function.

# 3) THE ADIPOCYTE AND ADIPOSE TISSUE<sup>[19]</sup>

Adipose tissue is composed of the lipidstoring adipose cell and a stromal/vascular compartment in which cells including preadipocytes and macrophages reside. Adipose mass increases by enlargement of adipose cells through lipid deposition, as well as by an increase in the number of adipocytes. Obese adipose tissue is also characterized by increased numbers of infiltrating macrophages. Although the adipocyte has generally been regarded as a storage depot for fat, it is also an endocrine cell that releases numerous molecules in a regulated fashion. These include the

energy balance regulating hormone leptin, cytokines such as tumor necrosis factor (TNF)- $\alpha$  and interleukin (IL)-6, complement factors such as factor D (also known as adipsin), prothrombotic agents such as plasminogen activator inhibitor I, and a component of the blood pressure–regulating system, angiotensinogen.

#### **G) MANAGEMENT OF OBESITY**

VaatghnanyannapananiShleshma-Medo-Harani Cha |

RukshoshnaBastaysteekshnaRukshanyudvartanani Cha  $||^{[20]}$ 

- 1} Charakacharya quotes for reducing the bulk of the obese heavy and non saturating therapy is prescribed. Food and drinks allevating Vata and reducing Kapha and Meda, Ruksha, Ushna and TeekshnaBasti, RukshaUdvartana (Rough Annointing), Use of Guduchi, Devdaru, Musta, Trifala, Takrarishta and Madhuis recommended in treatment of obesity.
- 2] Acharya Sushruta quoted use of Shilajatu, Guggulu, Gomutra, Trifala, Lohachurna, Rasanjana, Yava, Mudga, Koradusha, Shyamaka, UddalakaetcVirukshanakari, Chedana (causing depletion of Meda, Kapha) substances, appropriate use of LekhanaBasti, Vyayama. [21]
- 3} Ashtangasangrahakara Vagbhata quoted VataghnaAnnapana, Kapha-Medahara food substances like Kulattha, Yavaka, Yava, Jurna, Shyamaka, Mudga, Arishtapana, Madhudaka (Honey mixed with water), Mastu, Takra etc consumption, use of drugs having Teekshna, Ushna, Ruksha, ChchediGuna. Chinta, Vyayaya, Vyayama, Shodhana, Aswapana (night evil), Ruksha Snana, Udvartana in management of obesity<sup>[22]</sup>.

Modern Science focuses more precisely management of obesity it is as follows <sup>[23]</sup>,

#### 1) LIFESTYLE MANAGEMENT

Obesity care involves attention to three essential elements of lifestyle: dietary habits, physical activity, and behavior modification. Because obesity is fundamentally a disease of energy imbalance, all patients must learn how and when energy is consumed (diet), how and when energy is expended (physical activity), and how to incorporate this information into their daily lives (behavioral therapy). Lifestyle management has been shown to result in a modest (typically 3–5 kg) weight loss when compared with no treatment or usual care.

#### 2) DIET THERAPY

The primary focus of diet therapy is to reduce overall calorie consumption. The calorie deficit can be instituted through dietary substitutions. Examples include choosing smaller portion sizes, eating more fruits and vegetables, consuming more whole-grain cereals, selecting leaner cuts of meat and skimmed dairy products, reducing consumption of fried foods and other foods with added fats and oils, and drinking water instead of sugar-sweetened beverages. It is important that dietary counseling remain patient centered and that the goals set be practical, realistic, and achievable. The macronutrient composition of the diet will vary with the patient's preference and medical condition. The macronutrient composition will ultimately be determined by the patient's taste preferences, cooking style, and culture. Another dietary approach to consider is based on the concept of energy density, which refers to the number of calories (i.e., amount of energy)

a food contains per unit of weight. People tend to ingest a constant volume of food regardless of caloric or macronutrient content. Adding water or fibre to a food decreases its energy density by increasing weight without affecting caloric content. Examples of foods with low-energy density include soups, fruits, vegetables, oatmeal, and lean meats. Dry foods and high-fat foods such as pretzels, cheese, egg yolks, potato chips, and red meat have a high-energy density. Diets containing low-energy-dense foods have been shown to control hunger and thus to result in decreased caloric intake and weight loss.

## 3) PHYSICAL ACTIVITY THERAPY

Although exercise alone is only moderately effective for weight loss, the combination of dietary modification and exercise is the most effective behavioural approach for the treatment of obesity. The most important role of exercise appears to be in the maintenance of the weight loss. The 2008 Physical Activity Guidelines for Americans www.health.gov/paguidelinesrecommend that adults should engage in 150 min of moderateintensity or 75 min a week of vigorousintensity aerobic physical activity per week, performed in episodes of at least 10 min and preferably spread throughout the week. Focusing on simple ways to add physical activity into the normal daily routine through leisure activities, travel, and domestic work should be suggested. Examples include walking, using the stairs, doing housework and yard work, and engaging in sports. These exercise recommendations are daunting to most patients and need to be implemented gradually. Consultation with an exercise physiologist or personal trainer may be helpful.

#### 4) BEHAVIORAL THERAPY

Cognitive behavioural therapy is used to help change and reinforce new dietary and physical activity behaviours. Strategies include selfmonitoring techniques (e.g., journaling, weighing, and measuring food and activity); stress management; stimulus control (e.g., using smaller plates, not eating in front of the television or in the car); social support; problem solving; and cognitive restructuring to help patients develop more positive and realistic thoughts about themselves. When recommending any behavioural lifestyle change, the patient should be asked to identify what, when, where, and how the behavioural change will be performed. The patient should keep a record of the anticipated behavioural change so that progress can be reviewed at the next office visit. Because these techniques are timeconsuming to implement, their supervision is often undertaken by ancillary office staff, such as a nurse-clinician or registered dietician.

#### 5) PHARMACOTHERAPY

Adjuvant pharmacologic treatments should be considered for patients with a BMI ≥30 kg/m<sup>2</sup> or for patients who have concomitant obesityrelated diseases and for whom dietary and physical activity therapy has not been successful a BMI ≥27 kg/m2. Medications for obesity have traditionally fallen into two major categories: appetite suppressants (anorexiants) and fat gastrointestinal blockers. Appetitesuppressing medications have primarily targeted three monoamine receptor systems in the hypothalamus: noradrenergic, dopaminergic, and serotonergic receptors. Two new appetite suppressants were approved by the U.S. FDA in 2012: lorcaserin and phentermine/topiramate PHEN/TPM extended re-

lease. Gastrointestinal fat blockers reduce the absorption of selective macronutrients, such as fat, from the gastrointestinal tract. Among the anorexiants, phentermine has been the most commonly prescribed; there is limited long-term data on its effectiveness.

#### 6) SURGERY

Bariatric surgery can be considered for patients with severe obesity BMI,  $\geq$ 40 kg/m2 or for those with moderate obesity BMI,  $\geq$ 35 kg/m2 associated with a serious medical condition.

Weight loss surgeries have traditionally been classified into three categories on the basis of changes: restrictive, restictiveanatomic malabsorptive, and malabsorptive. More recently, however, the clinical benefits of bariatric surgery in achieving weight loss and alleviating metabolic comorbidities have been attributed largely to changes in the physiologic responses of gut hormones and in adipose tissue metabolism. The mortality rate from bariatric surgery is generally <1% but varies with the procedure, the patient's age and comorbid conditions, and the experience of the surgical team. The most common surgical complications include stomal stenosis or marginal ulcers occurring in 5-15% of patients that present as prolonged nausea and vomiting after eating or inability to advance the diet to solid foods. These complications typically are treated by endoscopic balloon dilation and acid suppression therapy, respectively.

## H) PATHYA- APATHYA<sup>[24]</sup>

#### A) Pathya in Sthaulya

Following substances are useful in Sthaulya

Anna varga, PuranaShali, Raktashali, Shasthikashali, Yava, Chanaka, Kulatha, ShrutaSheetaJala, PanchkolashrutaJala, Shunti siddha jala, Madhu, PuranaSidhu, Gomutra, Lasuna, Ardraka, Sunthi, Takra, Shakavarga, Patola, Karavellaka, Varthaka, Nimbapatra. Shigru, JangalaMamsa

#### B) Apathya in Sthaulya

Following substances are to be prohibited in Sthaulya

Masha, Taila, DushitaJala, SheetaJala, NutanaMadya, Aluka, Dadhi, Ksheera, GudaAnupaMamsa, Ahara having Guru Snigdha, Ati Drava, Picchila and AbhishyandiGuna

#### REFERENCES

- 1. https://en.wikipedia.org/wiki/Obesity#Epi demiology
- https://en.wikipedia.org/wiki/Obesity\_in\_I ndia
- Timesofindia.indiatimes.com/india/No-1in-underweight-population-India-amongtop-5-inobesity/articleshow/51656193.cms
- 4. Harrisons Principles of Internal Medicine written by Kasper, Fauci, Hauser, Longo, Jameson, Loscalzo 19<sup>th</sup> Edition Published by McGrawHill Education, Newyork2015 Page no 415e-1
- Sanskrit English Dictionary written by Sir Monier Williams Published by MotilalBanarasidass Publication New Delhi Reprint 1986 Page no.1266
- 6. https://en.wikipedia.org/wiki/Obesity#ety mology
- 7. Charaka Samhita by Agnivesha, revised by Charaka and Dridhabala with Ayurveda-Dipika commentary of Chakrapani-Datta

- edited by VaidyaY.G.JoshiSutrasthana 21<sup>st</sup>AdhyayaShloka no. 9 Published by Vaidyamitra Publication, Pune 2003
- 8. Charaka Samhita by Agnivesha, revised by Charaka and Dridhabala with Ayurveda-Dipika commentary of Chakrapani-Datta edited by VaidyaY.G.JoshiSutrasthana 23<sup>rd</sup> AdhyayaShloka no. 3,4 Published by Vaidyamitra Publication, Pune 2003
- Ashtangsangraham by Vruddha Vagbhata with Shashilekha Sanskrit Commentary of IndhuSutrasthanaAdhyaya 24<sup>th</sup> Page no.216 Edited by VaidyaGovardhan Sharma Changani Published by Choukhamba Sanskrit Series, Banaras 1945 Reprint 2010
- 10. Bhavprakash by Bhavmishrawith VidyotiniTeeka of Bramha Shankar Mishra Shastreeedited by VaidyaHarihar Prasad Pandey Madhya Khanda Chapter No.39 Shloka No.3-5 Published by Choukhamba Sanskrit Series Banaras 1998
- 11. https://en.wikipedia.org/wiki/Obesity# Causes
- 12. Charaka Samhita by Agnivesha, revised by Charaka and Dridhabala with Ayurveda-Dipika commentary of Chakrapani-Datta edited by VaidyaY.G.JoshiSutrasthana 21<sup>st</sup>AdhyayaShloka no. 4 Published by Vaidyamitra Publication, Pune 2003
- 13. Ashtangsangraham by Vruddha Vagbhata with Shashilekha Sanskrit Commentary of IndhuSutrasthanaAdhyaya 24<sup>th</sup> Page no.216 Edited by VaidyaGovardhan Sharma Changani Published by Choukhamba Sanskrit Series, Banaras 1945 Reprint 2010
- 14. Harrisons Principles of Internal Medicine written by Kasper, Fauci, Hauser, Longo, Jameson, Loscalzo 19<sup>th</sup> Edition Published

- by McGrawHill Education, Newyork 2015 Page no. 2394
- 15. Charaka Samhita by Agnivesha, revised by Charaka and Dridhabala with Ayurveda-Dipika commentary of Chakrapani-Datta edited by VaidyaY.G.JoshiSutrasthana 21<sup>st</sup>AdhyayaShloka no.5-8 Published by Vaidyamitra Publication, Pune 2003
- 16. Kayachikitsa written by VaidyaY.G.Joshi published by Pune SahityaVitarana Publication 2010 Chapter 21 Page no.212
- 17. https://en.wikipedia.org/wiki/Obesity# Pathology
- 18. Harrisons Principles of Internal Medicine written by Kasper, Fauci, Hauser, Longo, Jameson, Loscalzo 19<sup>th</sup> Edition Published by McGrawHill Education, Newyork 2015 Page no 415e-2
- 19. Harrisons Principles of Internal Medicine written by Kasper, Fauci, Hauser, Longo, Jameson, Loscalzo 19<sup>th</sup> Edition Published by McGrawHill Education, Newyork 2015 Page no 415e-2
- 20. Charaka Samhita by Agnivesha, revised by Charaka and Dridhabala with Ayurveda-Dipika commentary of Chakrapani-Datta edited by VaidyaY.G.JoshiSutrasthana 21<sup>st</sup>AdhyayaShloka no.21 Published by Vaidyamitra Publication, Pune 2003
- 21. Sushruta Samhita by Sushruta edited by VaidyaY.T.AcharyaSutrasthana chapter 15<sup>th</sup>Shloka no. 32 published by NirnayaSagar Press Bombay 1945
- 22. Ashtangsangraham by Vruddha Vagbhata with Shashilekha Sanskrit Commentary of IndhuSutrasthanaAdhyaya 24<sup>th</sup> Page no.217 Edited by VaidyaGovardhan Sharma Changani Published by Choukhamba Sanskrit Series, Banaras 1945 Reprint 2010

- 23. Harrisons Principles of Internal Medicine written by Kasper, Fauci, Hauser, Longo, Jameson, Loscalzo 19<sup>th</sup> Edition Published by McGrawHill Education, Newyork 2015 Page no 2394 to 2397
- 24. Kayachikitsa written by VaidyaY.G.Joshi published by Pune SahityaVitarana Publication 2010 Chapter 21 Page no.212

# Source of Support: Nil Conflict Of Interest: None Declared

How to cite this URL: Pranit Hanumantrao Patil: Critical Study Of Sthaulya (Obesity) And Its Principles Of Management. International Ayurvedic Medical Journal {online} 2017 {cited November, 2017} Available from:

http://www.iamj.in/posts/images/upload/4264 4276.pd