

PRILIMINARY PHYTOCHEMICAL SCREENING OF BASELLA RUBRA LINN

Manju Singh¹, Rajesh Kumari², Mita Kotecha³¹Women Scientist, Department of Biochemical Engineering and Biotechnology, IIT Delhi, Hauz Khas, New Delhi-110016,²Assistant Professor, Shiv Shakti Ayurvedic College, Bhikhi (Mansa)³Professor, H.O.D, Department of Dravyaguna Vigyan, National Institute of Ayurveda, Madhav Vilas, Amer Road, Jaipur.

ABSTRACT

Basella rubra linn belongs to family Basellaceae. The present paper includes macroscopy, preliminary phytochemical and physicochemical evaluation & TLC profile. Physicochemical parameters such as total ash value, acid insoluble ash value and water soluble ash value were determined which were 27.192 ± 1.171 % w/w, 0.7376 ± 0.2459 % w/w, 15.464 ± 1.879 % w/w respectively. Preliminary phytochemical analyses of extracts were carried out. The results were positive for carbohydrates, reducing sugar, protein, saponin, etc. These secondary metabolites are the active constituents of and it may be responsible for its pharmacological activities.

Keywords: Basella rubra, upodika, phytochemical and physicochemical

INTRODUCTION

Basella rubra L. (Basellaceae), commonly known as Indian or Malabar spinach belongs to family Basellaceae¹, is an herbaceous annual or biennial climbing herb found in tropical and sub-tropical areas. It is a succulent, branched, smooth, twining herbaceous vine, several meters in length. Stems are purplish or green. Leaves are fleshy, ovate or heart-shaped, 5 to 12 cms long, stalked, tapering to a pointed tip with a cordate base. Spikes are axillary, solitary, 5-29 cm long. Fruit is fleshy, stalkless, ovoid or spherical, 5-6 mm long, and purple when mature and contain only one seed. The flowers are pink, and about 4 millimeters long. The leaves of the plant contain flavonoids (133.1 ± 26.2 mg QC /100 g FM), β -cyanin

and 7, 4'-di-ortho methyl kempferol. The flower contain phenolic compounds (269.0 ± 3.1 mg GAE/100 g FM) such as Rutin, Quercetin, Scopoletin, Coumarin, β -xanthin and β -cyanin pigments and Caffeic-, Homo-protocatechuic-, Chlorogenic-, trans- and cis-p-coumaric-, p-hydroxybenzoic-, phloretic-, trans- and cis-sinapic-, cinnamic- acids; and the fruit consists of β -cyanin, gomphrenin I, gomphrenin II, and gomphrenin III². Basella rubra Linn. is a rich source of nutrients and minerals. Per 100 grams (g) edible portion, alugbati leaves contain Water (g) – 92.5; Energy (kcal) – 23.0; Protein (g) – 2.0; Fat (g) – 0.3; Carbohydrates (g) – 3.0; Fiber (g) – 0.9; Ash (g) – 2.2; Calcium (mg) – 128.0; Phosphorous (mg) – 40.0; Iron (mg) – 4.9;

Vitamin A (ug) – 456.0; Thiamine (mg) – 0.04; Riboflavin (mg) – 0.12; Niacin (mg) – 0.5; Ascorbic acid (mg) – 89.0²⁷. It also contains calcium 2.32, potassium 5.8, and magnesium 0.06, sodium 5.11 iron 0.04mg/100gm 3.

MATERIAL AND METHODS

Basella rubra leaves were collected from surata and Vadodara, Gujrata herbal garden and authenticated at Botanical Survey of India, Arid Zone regional circle, Jodhpur-08 (Rajasthan) on date 7th March 2008. Voucher specimens were collected and placed in the herbarium of Department of Dravyaguna vigyan, National Institute of Ayurveda. Prior to all analysis of all the raw materials were cleaned to remove any foreign materials and dust. The *Basella rubra linn*. leaves were dried in shade and powdered in electric grinder for physico-chemical, phytochemical and TLC studies. Prepared powdered was stored in an airtight container. For accuracy, each analysis was carried out in triplicate and average values of the parameters are reported. Plant images are shown in plate No1, 2 and 3. The various parameters analysed are as under:

ORGANO-LEPTIC & MACROSCOPICAL EVALUATION

The freshly grinded seed powder was evenly spread on a clean dry plastic sheet and Investigated for different organoleptic features by repeated observations by conventional Methods were recorded.

PHYSICO-CHEMICAL AND PHYTO-CHEMICAL PARAMETERS

Physicochemical and phytochemical screening for were carried out as per the guidelines given in Pharmacopoeia of India 2006⁴ Physico-chemical parameters such as moisture content (loss on drying at 105 C) , water soluble extractive value, alcohol soluble extractive value, total ash value, acid soluble ash value and water

soluble ash were calculated. The organic substance of the *Basella rubra linn* leaves show their solubility in various, solvents in different quantities. So for this purpose of determination of extractive values seven solvents (Benzene, Toluene, DCM, Ethyl acetate, Acetone, methanol and water) were selected according to their polarity. Coarsely powdered air dried drug material is accurately weighed and taken in a glass stopper conical flask. Solvent is added to the flask and the flask is attached to a reflux condense and boiled for 6 hours, on water bath. After 6 hours, the flask is allowed to cool and the content is filtered through filter paper. The filtrate is transferred to a pre- weighed flat bottomed dish and evaporated to dryness on a water bath. Then the dish kept in oven for six hours for the contents to get dried fully. The dish is cooled by keeping in a desiccator for 30 minutes and weighed without delay. The residual mass remained in filter paper is dried as such and is collected fully. This mass in again put into the conical flask and added with next solvent according to polarity, and fitted with reflux condenser, and extract is prepared in the same method used above. This procedure is repeated with all the seven solvents. The content of the extractable matter is calculated in mg per gms of air dried material. Preliminary tests were carried out on ethanolic and water extract for presence/ absence of phyto-constituents like carbohydrates, reducing sugar, protein, saponin, alkaloid, starch, tannin, phenol and glycosides^{4,5}.

The ethanolic extract of drug was subjected to thin layer chromatographic analysis. T.L.C. plates precoated with 0.25 mm layer of silica gel GF 254 with fluorescent indicator, (Merck's) were used. It was cut in size of 8×2 cm. ethanolic extract was applied on the plate above 1 cm from base line. The separated spots were visualized

under UV 366 nm. The qualitative evaluation of the plate was done by determining the migratory behavior of the separated substances given in the form of Rf values⁶.

OBSERVATIONS

Ayurvedic pharmacodynamics⁷⁻¹⁰

Rasa -Madhura

Guṇa – , pichhala, Snigdha and sara

Virya - Shita

Vipaka- Madhura

Effect on dosha- Vātaghna, Pittaghna, Kaphakara action

Organoleptic Character^{7,8}

Sparsh (Touch)- *Snigdha (smooth)*

Rupa (Apperance) – *Dark green*

Rasa (Taste)- *Madhura*

Gandha (Smell)- No particular smell

Table No.1

Total ash	Acid insoluble ash	Water soluble ash
27.192 ±1.171	0.7376 ± 0.245	15.464 ±1.879

Table No. 2- Extractive values of leaves of *Basella rubra*

S.No.	Solvent	Percentage of Extract
	Benzene	.0084%
	Toluene	.000264%
	Dichloromethane	.0026%
	Ethylacetate	0.001116%
	Acetone	.00733%
	Methanol	.011522%
	Water	.0002%

Table No. : Observation of Qualitative analysis of Inorganic matter in *Leaves of Basella rubra*.

S.No.	Name of the minerals / electrolytes	<i>Basella rubra</i> linn. leaves
1.	Calcium	+
2.	Iron	+
3.	Phosphorus	+
4.	Potassium	–
5.	Sulphur	+
6.	Cobalt	-
7.	Copper	-
8.	Mercury	-
9.	Nickel	-
10	Zinc	–
11.	Silver	-
13.	Zinc	- ve

Table No:4. Observation of Qualitative analysis of Organic matter in *Leaves of Basella rubra*.

S. No.	Chemical constituent	Test Applied	Result
	Carbohydrates	Molisch's reagent	+

	Reducing sugar	Fehling solution	+
	Starch	Iodine solution	+
	Tannin	Vanillin solution	-
	Protein	Ninhydrin solution	+
	Saponin	Shaking with water	+
	Phenol	FeCl ₃ solution	+
	Glycoside	FeCl ₃ + H ₂ SO ₄ solution	-
	Alkaloid	Dragondroff's reagent	-

T.L.C. of *Basella rubra* Linn.**Ethanol Extract**

Mobile phase- Chloroform:methanol :hexane : Benzene ::5:1:3.5 :1.5

Visualization: Short Wave (254 nm) UV

Table no: 5.1 Showing T.L.C of the leaves of *Basella rubra* of Ethanol Extract.

Spot No.	Distance traveled by solvent	Distance traveled by solute	R _f Value
1.	5.9	.3.	0.05084
2	5.9	2.5	0.4237
3	5.9	2.3	0.389
4	5.9	3	0.5084

Aqueous Extract

Mobile phase-Methanol:Petroleum ether :: 1:1

Visualization::Short Wave (254 nm) UV

Table no:5.2 Showing T.L.C of the leaves of *Basella rubra* of Aqueous Extract.

Spot No.	Distance traveled by solvent	Distance traveled by solute	R _f Value
1.	4.9	.2	0.04
2.	4.9	.5	0.10
3	4.9	.9	0.18

RESULTS AND CONCLUSION:

The Ash values are given in table no 1. Different extractive values are given in Table no 2. Preliminary phytochemical tests for inorganic and organic matter are given in table no 2& table no.4. Details of TLC were given in table no.5.1 and table no-5.1

This study is in line with the quality parameters prescribed in Ayurvedic Pharmacopeia of India and also standards set by other international agencies. This work provides qualitative and quantitative standards for the identification of *Basella rubra linn.* and from this study it is concluded that phytochemical studies on *Ba-*

sella rubra linn will be highly useful in determining qualitative and quantitative standards which can ascertain the identity, quality and purity of this plant drug.

REFERENCES

1. Guarino, L. (Editor), 1997. Traditional African vegetables. Proceedings of the IPGRI international workshop on genetic resources of traditional vegetables in Africa: conservation and use, 29-31
2. Dr.Gyandendra Pandey. Dravyaguna Vijnana. Varanasi: Krishnadas Academy; 2003 vol (3). 728-32
3. Nirmala, S. Saroja and G. Gayathri Devi 2011. Antidiabetic Activity of *Ba-*

- sella rubra and its Relationship with the Antioxidant Property A. British Biotechnology Journal 1(1), 1-9
- Mukherjee PK, Quality Control of Herbal Drugs, Business Horizons Pharmaceutical publishers, New Delhi, 2002, 356-357
 - Kumari Rajesh, Kotecha Mita. Physico-chemical and nutritional evaluation of Yava (*Hordeum vulgare* linn.) and Jai (*Avena sativa* linn.); Int. Res. J. Pharm 2015; 6(1); 70-72
 - Mendham J, Denney RC, Barnes JD and Thomas MJK: Vogel's Textbook of Quantitative Chemical Analysis, Pearson Education Pvt. Ltd., Six Edition 2002, 256-57
 - Ethnobotany and Medicinal Plants of Indian Subcontinent (J.K. Mahe-swari) pg 239
 - Prof. K.R. Srikantha Murty 2008. Suśrut Samhitā Sutra Sthana (English) Chaukhamba Orientalia, Varanasi: 1(1); 89, 2; 54, 55, 86, 88
 - Prof. Priyavrat Sharma 2003 Charak Samhitā Sutra Sthana (English) , Chaukhamba Orientalia , Varanasi : 1; 119, 229,
 - Satya Narayana Sastri 2003 Charak Samhitā Sutra Sthana (Hindi), Chaukhamba Bharati Academy: 1; 58, 348, 572, 713
 - Satya Narayana Sastri 2003 Charak Samhitā Cikitsa Sthana (Hindi) Chaukhamba Bharati Academy , Varanasi, .2, 100, 779, 780, 789.

Plate No 1- *Basella rubra* linn. Leaves Plate no.2 Dry Leaves Plate no.3 Dry Leaves Powder



CORRESPONDING AUTHOR

Dr Manju Singh

Women Scientist,

Department of Biochemical Engineering
and Biotechnology,

IIT Delhi, Hauz Khas, New Delhi-110016,

Email: manjubams@gmail.com

Source of Support: Nil

Conflict of Interest: None Declared